

10/048196


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

BOX PCT/DO-EO

Commissioner for Patents
Washington, D.C. 20231

"Express Mail" mailing label number EL 823 533 225 US

I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 CFR § 1.10 on the date indicated above and addressed to: Commissioner of Patents, Washington, DC 20231, on January 23, 2002.


Susan C. Sheehan

NATIONAL STAGE APPLICATION TRANSMITTAL LETTER
APPLICATION FILING UNDER 35 U.S.C. § 371

Transmitted herewith for filing is the patent application of:

Inventor(s)/Applicant(s): Thonnard, Joelle et al.
International Application No.: EP00/07291
International Published Appln. No.: WO 01/09330
International Filing Date: 27 July 2000
Priority Filing Date: 30 July 1999
Thirty Month Date: 30 January 2002
Title: "Novel Compounds"

1. **THIS NEW APPLICATION IS A NATIONAL STAGE APPLICATION UNDER PCT, CHAPTER II WITH A REQUEST FOR EXAMINATION WITHOUT DELAY TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US).**

- ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. § 371;
☐ This is a **SECOND** or subsequent submission of items concerning a filing under 35 U.S.C. § 371.

2. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).

3. A proper Demand for International Preliminary Examination was made by the 19TH month from the earliest claimed priority date.
4. Enclosed items are required for filing under 37 CFR § 1.53(b) and § 1.494(b) or § 1.495(b):

- ☒ One copy of International Publication WO 01/09330
- (a) ☒ is transmitted herewith (**required only if not transmitted by the International Bureau**)
- (b) ☒ has been transmitted by the International Bureau
- (c) ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)

Fees

☒ The basic national fee set forth in 37 CFR § 1.482 - International Preliminary Examination Fee not paid to USPTO but International Search Report prepared by the EPO or JPO - **\$860.00**

- ☒ Claims in Excess of 20 (20 @ \$18.00)
- ☐ Independent Claims in Excess of 3 (4 @ \$80.00)

5. Further enclosed are:

- ☒ One copy of International Preliminary Examination Report.
- ☒ One copy of International Search Report - (with references).

Information Disclosure Statement:

- The Examiner is requested to consider carefully the complete text of the documents submitted herewith in connection with the examination of this application. It is believed that the Examiner will concur with Applicant's belief that the documents do not adversely affect the patentability of the subject matter presently claimed, taken alone or in combination.
- It is requested that the listed documents be included in the "References Cited" portion of any patent issuing from this application.
- Under 37 CFR 1.97(i), Applicants understand that non-complying Information Disclosure Statements will be placed in the file but not considered by the Office, however, under Reply to Comment 8 of the Federal Register, page 2024, Applicants will be informed when information is not considered.

[X] Applicants enclose and make of record the documents identified in the International Search Report or in the International Preliminary Examination Report. These documents are listed on the enclosed Forms PTO/SB/08A and PTO/SB/08B.

- ☐ One copy of Written Opinion.
- ☐ One copy of PCT Request as filed.
- ☐ One copy of Chapter II Demand as filed.

6. ☐ A translation of the International Application into English (35 U.S.C. § 371(c)(2))

7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. § 371(c)(3))

8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. § 371(c)(3))

9. Still additional papers enclosed:

- ☐ Assignment with Assignment Recordation Form Cover Sheet
- ☐ Verification Statement Claiming Small Entity Status
- ☒ Declaration or oath is enclosed executed by the inventor
- ☐ An Information Disclosure Statement under 37 CFR § 1.97 and § 1.98
- ☒ Return Acknowledgment Postcard

10. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)

A. Enclosed are:

- (a) ☒ Computer Readable Copy of the Sequence Listing
- (b) ☒ Paper Copy (identical to Computer Readable Copy) of the Sequence Listing

B. ☐ Enclosed is a paper copy of the Sequence Listing. This paper copy and a Computer Readable Form thereof are identical with the Computer Readable Form in another application of the Applicant which is fully identified as follows:

U.S. Application No.: @@

Filed: @@

Attorney Docket No.: @@

which is believed to comply with the rules set forth in 37 CFR § 1.821 et. seq. Applicants requests pursuant to 37 CFR § 1.821(e) that this Computer Readable Form be used in the present application. **Please TRANSFER the sequence listing from the parent to this application.**

C. ☒ Statement under 37 CFR § 1.821(f): **The information recorded in computer readable form is identical to the written Sequence Listing.**

D. ☒ Statement under 37 CFR § 1.821(g) (required when Sequence Listing not submitted at the time of filing under 35 U.S.C. §111(a)) or 37 CFR §1.821(f) (required when Sequence Listing not submitted at the time of

filing under the Patent Cooperation Treaty): The submission of the

Sequence Listing includes no new matter.

E. ☒ Amendment: Please enter the Sequence Listing into the application.

11. **Preliminary Amendment**

Prior to calculation of fees, kindly enter:

☒ Preliminary Amendment submitted herewith

☐ do not enter Preliminary Amendment

12. Fee payment being made at this time is enclosed:

*	Basic filing fee (\$860.00)	860.00
*	Claims in Excess of 20	54.00
*	(20 @ \$18.00)	
	Independent Claims in Excess of 3	.00
	(4 @ \$80.00)	
*	Total Fees enclosed:	<u>\$914.00</u>

13. The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Account No. 50-0258. This letter is filed in duplicate for accounting purposes.

Respectfully submitted,

Date: January 23, 2002



Teresa O. Bittenbender
Registration No. 47,425
Attorney for Applicant

DECHERT
4000 Bell Atlantic Tower
1717 Arch Street
Philadelphia, PA 19103-2793
Fax: (215) 994-2222
Tel: (215) 994-2213

International Application No.: PCT/EP00/07291
Attorney Docket No.: BM45402

#4/a

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Thonnard, et al.
Docket No.: BM45402
Serial No.: Unknown
Filed: Herewith
For: Novel Compounds

Group Art Unit No.: Unknown
Examiner: Unknown

PRELIMINARY AMENDMENT

Sir:

Applicant respectfully requests that this Preliminary Amendment be entered in this case before the calculation of fees and before examination of the subject application.

In the Claims:

Please delete the claims of the application as filed in the PCT and substitute therefor:

27. An isolated polypeptide comprising a member selected from the group consisting of
 - (a) an amino acid sequence matching SEQ ID NO:2 and
 - (b) an immunogenic polypeptide comprising a fragment sequence of at least 15 amino acids that matches an aligned contiguous segment of SEQ ID NO:2,wherein the isolated polypeptide, when administered to a subject in a suitable composition which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an antibody or T-cell immune response to a polypeptide having the sequence of SEQ ID NO:2.
28. An isolated polynucleotide encoding a polypeptide of Claim 27 or the full complement to the isolated polynucleotide.
29. The isolated polypeptide of claim 27, wherein the polypeptide is according to (a).
30. An isolated polynucleotide encoding a polypeptide of Claim 29 or the full complement to the isolated polynucleotide.
31. A process for expressing the polynucleotide of Claim 30 comprising transforming a host cell with an expression vector comprising the polynucleotide and culturing the host cell under conditions sufficient for expression of the polynucleotide.

32. The isolated polypeptide of claim 27, wherein the polypeptide is according to (b).
33. An isolated polynucleotide encoding a polypeptide of Claim 32 or the full complement to the isolated polynucleotide.
34. The isolated polypeptide of claim 27, wherein the immunogenic fragment of (b) comprises at least 20 amino acids.
35. The isolated polypeptide of claim 27, wherein the isolated polypeptide consists of SEQ ID NO:2.
36. An isolated polynucleotide encoding the polypeptide of Claim 35 or the full complement to the isolated polynucleotide.
37. A process for expressing the polynucleotide of Claim 36 comprising transforming a host cell with an expression vector comprising the polynucleotide and culturing the host cell under conditions sufficient for expression of the polynucleotide.
38. A fusion protein comprising the isolated polypeptide of Claim 21.
39. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1.
40. An isolated polynucleotide segment comprising a polynucleotide sequence or the full complement of the entire length of the polynucleotide sequence, wherein the polynucleotide sequence hybridizes to the full complement of SEQ ID NO:1 minus the complement of any stop codon, wherein the hybridization conditions include incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing in 0.1x SSC at 65°C; and, wherein the polynucleotide sequence is identical to SEQ ID NO:1 minus any terminal stop codon, except that, over the entire length corresponding to SEQ ID NO:1 minus any terminal

stop codon, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression

$$n_n \leq x_n - (x_n \cdot y)$$

wherein x_n is the total number of nucleotides in SEQ ID NO:1 minus any terminal stop codon, y is at least 0.95, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n ; and wherein the polynucleotide sequence detects *Moraxella catarrhalis*.

41. An expression vector comprising the isolated polynucleotide of Claim 28.
42. A host cell transformed with the expression vector of Claim 41.
43. A vaccine comprising the polypeptide of Claim 27 and a pharmaceutically acceptable carrier.
44. The vaccine of Claim 43, wherein the vaccine comprises at least one other *Moraxella catarrhalis* antigen.
45. An antibody immunospecific for the polypeptide or immunogenic fragment of Claim 27.
46. A method for inducing an immune response in a mammal comprising administration of the polypeptide of Claim 27.
47. A method of diagnosing a *Moraxella catarrhalis* infection, comprising identifying a polypeptide of Claim 27, or an antibody that is immunospecific for the polypeptide, present within a biological sample from an animal suspected of having such an infection.
48. A method for inducing an immune response in a mammal comprising administration of the isolated polynucleotide of Claim 28.

49. A therapeutic composition useful in treating humans with *Moraxella catarrhalis* comprising at least one antibody directed against the polypeptide of claim 27 and a suitable pharmaceutical carrier.

REMARKS

Claims

Claims 1-26 have been canceled without prejudice or disclaimer of the subject matter therein. Applicant reserves the right to prosecute, in one or more patent applications, the canceled claims, the claims as originally filed, and any other claims supported by the specification.

New claims 27-49 have been introduced. No new matter is added.

Support

Support for the new claims is either apparent, or is as described in the text below. Support for the hybridization conditions may be found, for example, at page 14, fourth full paragraph. Support for the recitation of sequence relatedness such as in claim 40 may be found in the specification, for example, at pages 46-50.

Closing Remarks

Allowance of the pending claims is respectfully requested.

Respectfully submitted,

Frank O. Butler

Teresa O. Bittenbender
Registration No. 47,425
Attorney for Applicant

DECHERT
1717 Arch Street
4000 Bell Atlantic Tower
Philadelphia, PA 19103
Fax: (215) 994-2222
Attn: Teresa O. Bittenbender, Esq.
(215) 994-2213

Novel Compounds

FIELD OF THE INVENTION

This invention relates to polynucleotides, (herein referred to as "BASB125 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB125" or "BASB125 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the invention relates to diagnostic assays for detecting infection of certain pathogens.

BACKGROUND OF THE INVENTION

Moraxella catarrhalis (also named *Branhamella catarrhalis*) is a Gram-negative bacteria frequently isolated from the human upper respiratory tract. It is responsible for several pathologies the main ones being otitis media in infants and children, and pneumonia in elderlies. It is also responsible of sinusitis, nosocomial infections and less frequently of invasive diseases.

Otitis media is an important childhood disease both by the number of cases and its potential sequelae. More than 3.5 millions cases are recorded every year in the United States, and it is estimated that 80 % of the children have experienced at least one episode of otitis before reaching the age of 3 (Klein, JO (1994) Clin.Inf.Dis 19:823). Left untreated, or becoming chronic, this disease may lead to hearing losses that could be temporary (in the case of fluid accumulation in the middle ear) or permanent (if the auditive nerve is damaged). In infants, such hearing losses may be responsible for a delayed speech learning.

Three bacterial species are primarily isolated from the middle ear of children with otitis media: *Streptococcus pneumoniae*, non typeable *Haemophilus influenzae* (NTHi) and *M. catarrhalis*. They are present in 60 to 90 % of the cases. A review of recent studies shows that *S. pneumoniae* and NTHi represent both about 30 %, and *M. catarrhalis* about 15 % of the otitis media cases (Murphy, TF (1996) Microbiol.Rev. 60:267). Other bacteria could be

isolated from the middle ear (*H. influenzae* type B, *S. pyogenes* etc) but at a much lower frequency (2 % of the cases or less).

5 Epidemiological data indicate that, for the pathogens found in the middle ear, the
colonization of the upper respiratory tract is an absolute prerequisite for the development of
an otitis; other are however also required to lead to the disease (Dickinson, DP et al. (1988)
J. Infect.Dis. 158:205, Faden, HL et al. (1991) Ann.Otorhinol.Laryngol. 100:612). These are
important to trigger the migration of the bacteria into the middle ear via the Eustachian
tubes, followed by the initiation of an inflammatory process. These factors are unknown
10 todote. It has been postulated that a transient anomaly of the immune system following a
viral infection, for example, could cause an inability to control the colonization of the
respiratory tract (Faden, HL et al (1994) J. Infect.Dis. 169:1312). An alternative explanation
is that the exposure to environmental factors allow a more important colonization of some
children, who subsequently become susceptible to the development of otitis media because
15 of the sustained presence of middle ear pathogens (Murphy, TF (1996) Microbiol.Rev.
60:267).

The immune response to *M. catarrhalis* is poorly characterized. The analysis of strains
isolated sequentially from the nasopharynx of babies followed from 0 to 2 years of age,
20 indicates that they get and eliminate frequently new strains. This indicates that an efficacious
immune response against this bacteria is mounted by the colonized children (Faden, HL et al
(1994) J. Infect.Dis. 169:1312).

In most adults tested, bactericidal antibodies have been identified (Chapman, AJ et al. (1985)
25 J. Infect.Dis. 151:878). Strains of *M. catarrhalis* present variations in their capacity to resist
serum bactericidal activity: in general, isolates from diseased individuals are more resistant
than those who are simply colonized (Hol, C et al. (1993) Lancet 341:1281, Jordan, KL et al.
(1990) Am.J.Med. 88 (suppl. 5A):28S). Serum resistance could therefore be considered as a
virulence factor of the bacteria. An opsonizing activity has been observed in the sera of

children recovering from otitis media.

The antigens targetted by these different immune responses in humans have not been identified, with the exception of OMP B1, a 84 kDa protein which expression is regulated by iron, and that is recognized by the sera of patients with pneumonia (Sethi, S, et al. (1995) Infect.Immun. 63:1516), and of UspA1 and UspA2 (Chen D. et al.(1999), Infect.Immun. 67:1310).

A few other membrane proteins present on the surface of *M. catarrhalis* have been characterized using biochemical method, or for their potential implication in the induction of a protective immunity (for review, see Murphy, TF (1996) Microbiol.Rev. 60:267). In a mouse pneumonia model, the presence of antibodies raised against some of them (UspA, CopB) favors a faster clearance of the pulmonary infection. Another polypeptide (OMP CD) is highly conserved among *M. catarrhalis* strains, and presents homologies with a porin of *Pseudomonas aeruginosa*, which has been demonstrated efficacious against this bacterium in animal models.

The frequency of *Moraxella catarrhalis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Moraxella catarrhalis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

SUMMARY OF THE INVENTION

The present invention relates to BASB125, in particular BASB125 polypeptides and BASB125 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and

polynucleotides, including prevention and treatment of microbial diseases, amongst others. In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting expression or activity of BASB125 polynucleotides or polypeptides.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to BASB125 polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of BASB125 of *Moraxella catarrhalis*, which is related by amino acid sequence homology to none of the known proteins. It is predicted to be a lipoprotein because it has a signal sequence characteristic of lipoprotein. The invention relates especially to BASB125 having the nucleotide and amino acid sequences set out in SEQ ID NO:1 and SEQ ID NO:2 respectively. It is understood that sequences recited in the Sequence Listing below as "DNA" represent an exemplification of one embodiment of the invention, since those of ordinary skill will recognize that such sequences can be usefully employed in polynucleotides in general, including ribopolynucleotides.

Polypeptides

In one aspect of the invention there are provided polypeptides of *Moraxella catarrhalis* referred to herein as "BASB125" and "BASB125 polypeptides" as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

The present invention further provides for:

(a) an isolated polypeptide which comprises an amino acid sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% or exact identity, to that of SEQ ID NO:2;

(b) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1 over the entire length of SEQ ID NO:1; or

(c) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity, to the amino acid sequence of SEQ ID NO:2.

The BASB125 polypeptides provided in SEQ ID NO:2 is the BASB125 polypeptide from *Moraxella catarrhalis* strain MC2931 (ATCC 43617).

The invention also provides an immunogenic fragment of a BASB125 polypeptide, that is, a contiguous portion of the BASB125 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2; That is to say, the fragment (if necessary when coupled to a carrier) is capable of raising an immune response which recognises the BASB125 polypeptide. Such an immunogenic fragment may include, for example, the BASB125 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB125 according to the invention comprises substantially all of the extracellular domain of a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 over the entire length of SEQ ID NO:2

A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with

BASB125 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

5 Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2 or of a variant thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes
10 such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

15 Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or
20 deleted from the amino acid sequence of SEQ ID NO:2.

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments may be employed as intermediates for producing the full-length polypeptides of the invention.

25 Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

The polypeptides, or immunogenic fragments, of the invention may be in the form of
30 the "mature" protein or may be a part of a larger protein such as a precursor or a fusion

protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared to non-fused protein. The fusion partner may assist in providing T helper epitopes (immunological fusion partner), preferably T helper epitopes recognised by humans, or assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

Fusion partners include protein D from *Haemophilus influenzae* and the non-structural protein from influenza virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesize an N-acetyl-L-alanine
5 amidase LytA, (coded by the *lytA* gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E.coli* C- LytA expressing plasmids useful for expression of fusion
10 proteins. Purification of hybrid proteins containing the C- LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188 - 305.

15 The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or
20 aromatic residues Phe and Tyr.

Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a
25 combination of these methods. Means for preparing such polypeptides are well understood in the art.

It is most preferred that a polypeptide of the invention is derived from *Moraxella catarrhalis*, however, it may preferably be obtained from other organisms of the same

taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

Polynucleotides

5 It is an object of the invention to provide polynucleotides that encode BASB125 polypeptides, particularly polynucleotides that encode the polypeptide herein designated BASB125.

10 In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB125 polypeptides comprising a sequence set out in SEQ ID NO:1 which includes a full length gene, or a variant thereof.

The BASB125 polynucleotide provided in SEQ ID NO:1 is the BASB125 polynucleotide from *Moraxella catarrhalis* strain MC2931 (ATCC 43617).

15 As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB125 polypeptides and polynucleotides, particularly *Moraxella catarrhalis* BASB125 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B-
20 and Z-DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

Another aspect of the invention relates to isolated polynucleotides, including at least one full
25 length gene, that encodes a BASB125 polypeptide having a deduced amino acid sequence of SEQ ID NO:2 and polynucleotides closely related thereto and variants thereof.

In another particularly preferred embodiment of the invention there is a BASB125 polypeptide from *Moraxella catarrhalis* comprising or consisting of an amino acid
30 sequence of SEQ ID NO:2 or a variant thereof.

Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID NO:1, a polynucleotide of the invention encoding BASB125 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Moraxella catarrhalis* Catlin cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1, typically a library of clones of chromosomal DNA of *Moraxella catarrhalis* Catlin in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene sequence. Illustrative of the invention, the polynucleotide set out in SEQ ID NO:1 was discovered in a DNA library derived from *Moraxella catarrhalis*.

Moreover, the DNA sequence set out in SEQ ID NO:1 contains an open reading frame encoding a protein having about the number of amino acid residues set forth in SEQ ID NO:2 with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art.

The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 403 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

- 5 In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:
- (a) a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1 over the entire length of SEQ ID NO:1; or
 - 10 (b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID NO:2.
- 15 A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Moraxella catarrhalis*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions (for example, using a temperature in the range of 45 – 65°C and an SDS concentration from 0.1 – 1%) with a labeled or detectable probe consisting of or comprising
- 20 the sequence of SEQ ID NO:1 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

- The invention provides a polynucleotide sequence identical over its entire length to a coding sequence (open reading frame) in SEQ ID NO:1. Also provided by the invention is a coding
- 25 sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5'
- 30 and 3' sequence, such as the transcribed but non-translated sequences, termination signals

(such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson *et al.*, *Cell* 37: 767 (1984), both of which may be useful in purifying polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

The nucleotide sequence encoding BASB125 polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in nucleotides 1 to 402 of SEQ ID NO:1. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Moraxella catarrhalis* BASB125 having an amino acid sequence set out in SEQ ID NO:2. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2.

Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding BASB125 variants, that have the amino acid sequence of BASB125 polypeptide of SEQ ID NO:2 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of BASB125 polypeptide.

Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB125 polypeptide having an amino acid sequence set out in SEQ ID NO:2, and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 90% identical over its entire length to a polynucleotide encoding BASB125 polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 95% identical over their entire length to the same are particularly preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of SEQ ID NO:1.

In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB125 polynucleotide sequences, such as the polynucleotides in SEQ ID NO:1.

The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB125 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB125 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or

base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

5 A coding region of a BASB125 gene may be isolated by screening using a DNA sequence provided in SEQ ID NO:1 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

10 There are several methods available and well known to those skilled in the art to obtain full-length DNAs, or extend short DNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, *et al.*, *PNAS USA* 85: 8998-9002. 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly
15 simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that
20 is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a
25 separate full-length PCR using the new sequence information for the design of the 5' primer.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases,
30 particularly human diseases, as further discussed herein relating to polynucleotide assays.

The polynucleotides of the invention that are oligonucleotides derived from a sequence of SEQ ID NO:1 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in

combination with adjacent nucleotide positions, when read in the correct reading frame. would have the effect of generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein. a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein. or a preproprotein. which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff *et al.*, *Hum Mol Genet* (1992) 1: 363, Manthorpe *et al.*, *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu *et al.*, *J Biol Chem.* (1989) 264: 16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda *et al.*, *Science* (1989) 243: 375), particle bombardment (Tang *et al.*, *Nature* (1992) 356:152, Eisenbraun *et al.*, *DNA Cell Biol* (1993) 12: 791) and *in vivo* infection using cloned retroviral vectors (Seeger *et al.*, *PNAS USA* (1984) 81: 5849).

Vectors, Host Cells, Expression Systems

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation

systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis, *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, *Neisseria meningitidis* and *Moraxella catarrhalis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal
5 elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender
10 expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

15 In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be
20 heterologous signals.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose
25 chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), *Listeria*, *Salmonella*, *Shigella*, *Neisseria*, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

Diagnostic, Prognostic, Serotyping and Mutation Assays

This invention is also related to the use of BASB125 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB125 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB125 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from

a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled BASB125 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers *et al.*, *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for example, Cotton *et al.*, *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

In another embodiment, an array of oligonucleotide probes comprising BASB125 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee *et al.*, *Science*, 274: 610 (1996)).

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:
(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1, or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

(c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a Disease, among others.

- 5 This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferable, SEQ ID NO:1, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease, which results
- 10 from under-expression, over-expression or altered expression of the polynucleotide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.
- 15 Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such
- 20 as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding BASB125 polypeptide can be used to identify and analyze mutations.

- The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5'
- 25 and/or the 3' end. These primers may be used for, among other things, amplifying BASB125 DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide

sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections caused by *Moraxella catarrhalis*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of SEQ ID NO:1. Increased or decreased expression of a BASB125 polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of BASB125 polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a BASB125 polypeptide, in a sample derived from a host, such as a bodily material, are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probe obtained or derived from a bodily sample, to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Moraxella catarrhalis*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide

sequence of SEQ ID NO:1 are preferred. Also preferred is a grid comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2.

5 **Antibodies**

The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their
10 affinity for other related polypeptides in the prior art.

In certain preferred embodiments of the invention there are provided antibodies against BASB125 polypeptides or polynucleotides.

15 Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced
20 by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

25 Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

30

Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-BASB125 or from naive libraries (McCafferty, *et al.*, (1990), *Nature* 348, 552-554; Marks, *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson *et al.*, (1991) *Nature* 352: 628).

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

Thus, among others, antibodies against BASB125-polypeptide or BASB125-polynucleotide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

Antagonists and Agonists - Assays and Molecules

Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural

substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan *et al.*, *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

5 The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB125 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB125 polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and BASB125 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and and/or functionally related polypeptides (see D. Bennett *et al.*, *J Mol Recognition*, 8:52-58 (1995); and K. Johanson *et al.*, *J Biol Chem*, 270(16):9459-9471 (1995)).

30 The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for

detecting the effect of added compounds on the production of mRNA and/or polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or
5 enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of BASB125 polypeptides or
10 polynucleotides, particularly those compounds that are bacteriostatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB125 polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence
15 or the presence of a candidate molecule that may be a BASB125 agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the BASB125 polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of BASB125 polypeptide are most likely to be good antagonists. Molecules that bind well and,
20 as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate
25 converted into product, a reporter gene that is responsive to changes in BASB125 polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for BASB125 agonists is a competitive assay that combines BASB125 and a potential agonist with BASB125-binding molecules, recombinant
30 BASB125 binding molecules, natural substrates or ligands, or substrate or ligand mimetics,

under appropriate conditions for a competitive inhibition assay. BASB125 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB125 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

5

Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB125-induced activities, thereby preventing the action or expression of BASB125 polypeptides and/or polynucleotides by excluding BASB125 polypeptides and/or polynucleotides from binding.

10

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB125.

20

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

25

30

Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB125 proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

In accordance with yet another aspect of the invention, there are provided BASB125 agonists and antagonists, preferably bacteriostatic or bactericidal agonists and antagonists.

The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

In a further aspect, the present invention relates to mimotopes of the polypeptide of the invention. A mimotope is a peptide sequence, sufficiently similar to the native peptide (sequentially or structurally), which is capable of being recognised by antibodies which
5 recognise the native peptide; or is capable of raising antibodies which recognise the native peptide when coupled to a suitable carrier.

Peptide mimotopes may be designed for a particular purpose by addition, deletion or substitution of elected amino acids. Thus, the peptides may be modified for the
10 purposes of ease of conjugation to a protein carrier. For example, it may be desirable for some chemical conjugation methods to include a terminal cysteine. In addition it may be desirable for peptides conjugated to a protein carrier to include a hydrophobic terminus distal from the conjugated terminus of the peptide, such that the free unconjugated end of the peptide remains associated with the surface of the carrier
15 protein. Thereby presenting the peptide in a conformation which most closely resembles that of the peptide as found in the context of the whole native molecule. For example, the peptides may be altered to have an N-terminal cysteine and a C-terminal hydrophobic amidated tail. Alternatively, the addition or substitution of a D-stereoisomer form of one or more of the amino acids may be performed to create a
20 beneficial derivative, for example to enhance stability of the peptide.

Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number
25 of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB125 polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Moraxella catarrhalis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of BASB125 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB125 polynucleotide and/or polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB125 polynucleotide and/or polypeptide encoded therefrom, wherein the composition comprises a recombinant BASB125 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB125 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

A BASB125 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

In a vaccine composition according to the invention, a BASB125 polypeptide and/or polynucleotide, or a fragment, or a mimotope, or a variant thereof may be present in a vector, such as the live recombinant vectors described above for example live bacterial vectors.

Also suitable are non-live vectors for the BASB125 polypeptide, for example bacterial outer-membrane vesicles or "blebs". OM blebs are derived from the outer membrane of the two-layer membrane of Gram-negative bacteria and have been documented in many Gram-negative bacteria (Zhou, L *et al.* 1998. *FEMS Microbiol. Lett.* 163:223-228) including *C. trachomatis* and *C. psittaci*. A non-exhaustive list of bacterial pathogens reported to produce blebs also includes: *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella melitensis*, *Brucella ovis*, *Escherichia coli*, *Haemophilus influenza*, *Legionella pneumophila*, *Moraxella catarrhalis*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa* and *Yersinia enterocolitica*.

Blebs have the advantage of providing outer-membrane proteins in their native conformation and are thus particularly useful for vaccines. Blebs can also be improved

for vaccine use by engineering the bacterium so as to modify the expression of one or more molecules at the outer membrane. Thus for example the expression of a desired immunogenic protein at the outer membrane, such as the BASB125 polypeptide, can be introduced or upregulated (e.g. by altering the promoter). Instead or in addition, the
5 expression of outer-membrane molecules which are either not relevant (e.g. unprotective antigens or immunodominant but variable proteins) or detrimental (e.g. toxic molecules such as LPS, or potential inducers of an autoimmune response) can be downregulated. These approaches are discussed in more detail below.

10 The non-coding flanking regions of the BASB125 gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as
15 the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the shine-dalgarno sequence. regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences. This sequence is a further aspect of the invention.

20 This sequence information allows the modulation of the natural expression of the BASB125 gene. The upregulation of the gene expression may be accomplished by altering the promoter, the shine-dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modification. Alternatively, by changing phase variation
25 sequences, the expression of the gene can be put under phase variation control, or it may be uncoupled from this regulation. In another approach, the expression of the gene can be put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation include, but are not limited to, induction by temperature shift, addition of inductor substrates like selected carbohydrates or their derivatives. trace
30 elements, vitamins, co-factors, metal ions, etc.

Such modifications as described above can be introduced by several different means. The modification of sequences involved in gene expression can be carried out *in vivo* by random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating the region of interest and modifying it by random mutagenesis, or site-directed replacement, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the wild-type regulatory sequences. These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome. A non-exhaustive list of preferred promoters that could be used for up-regulation of gene expression includes the promoters *porA*, *porB*, *lbpB*, *tbpB*, *p110*, *lst*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*; *ompCD*, *copB*, *lbpB*, *ompE*, *UspA1*; *UspA2*; *TbpB* from *M. Catarrhalis*; *p1*, *p2*, *p4*, *p5*, *p6*, *lpD*, *tbpB*, *D15*, *Hia*, *Hmw1*, *Hmw2* from *H. influenzae*.

In one example, the expression of the gene can be modulated by exchanging its promoter with a stronger promoter (through isolating the upstream sequence of the gene, *in vitro* modification of this sequence, and reintroduction into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium.

In other examples, the described approaches can be used to generate recombinant bacterial strains with improved characteristics for vaccine applications. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the

immune response, strains with modulated expression of immunodominant proteins, strains with modulated shedding of outer-membrane vesicles.

Thus, also provided by the invention is a modified upstream region of the BASB125 gene, which modified upstream region contains a heterologous regulatory element which alters the expression level of the BASB125 protein located at the outer membrane. The upstream region according to this aspect of the invention includes the sequence upstream of the BASB125 gene. The upstream region starts immediately upstream of the BASB125 gene and continues usually to a position no more than about 1000 bp upstream of the gene from the ATG start codon. In the case of a gene located in a polycistronic sequence (operon) the upstream region can start immediately preceding the gene of interest, or preceding the first gene in the operon. Preferably, a modified upstream region according to this aspect of the invention contains a heterologous promotor at a position between 500 and 700 bp upstream of the ATG.

Thus, the invention provides a BASB125 polypeptide, in a modified bacterial bleb. The invention further provides modified host cells capable of producing the non-live membrane-based bleb vectors. The invention further provides nucleic acid vectors comprising the BASB125 gene having a modified upstream region containing a heterologous regulatory element.

Further provided by the invention are processes to prepare the host cells and bacterial blebs according to the invention.

Also provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Moraxella catarrhalis*.

5 Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial
10 infection, particularly *Moraxella catarrhalis* infection, in mammals, particularly humans.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and
15 polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostatic compounds and solutes which render the formulation isotonic with
20 the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

25

The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (*Mosmann, T.R. and Coffman, R.L. (1989) TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology, 7, p145-173*). Traditionally, TH1-type responses are associated with the production of the INF- γ and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2-type responses are associated with the secretion of IL-4, IL-5, IL-6 and IL-13.

It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2 - type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes
5 direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes *in vitro* after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell
10 populations to produce high levels of TH1-type cytokines when re-stimulated with antigen *in vitro*, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

Adjuvants which are capable of preferential stimulation of the TH1 cell response are
15 described in International Patent Application No. WO 94/00153 and WO 95/17209.

3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi
20 Immunochem, Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a 0.22micron membrane (European Patent number 0 689 454).
25 3D-MPL will be present in the range of 10µg - 100µg preferably 25-50µg per dose wherein the antigen will typically be present in a range 2-50µg per dose.

Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of Quillaja Saponaria Molina. Optionally this may be admixed

with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with an carrier.

The method of production of QS21 is disclosed in US patent No. 5,057,540.

Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; preferably 1:5 to 5 : 1 and often substantially 1 : 1. The preferred range for optimal synergy is 2.5 : 1 to 1 : 1 3D-MPL: QS21.

Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1 µg - 200 µg, such as 10-100 µg, preferably 10 µg - 50 µg per dose.

Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha
5 tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene: alpha
tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85
may also be present at a level of 1%. In some cases it may be advantageous that the
vaccines of the present invention will further contain a stabiliser.

10 Non-toxic oil in water emulsions preferably contain a non-toxic oil. e.g. squalane or
squalene, an emulsifier, e.g. Tween 80, in an aqueous carrier. The aqueous carrier may
be, for example, phosphate buffered saline.

A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol
15 in an oil in water emulsion is described in WO 95/17210.

The present invention also provides a polyvalent vaccine composition comprising a
vaccine formulation of the invention in combination with other antigens, in particular
antigens useful for treating cancers, autoimmune diseases and related conditions. Such a
20 polyvalent vaccine composition may include a TH-1 inducing adjuvant as hereinbefore
described.

While the invention has been described with reference to certain BASB125 polypeptides
and polynucleotides, it is to be understood that this covers fragments of the naturally
25 occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides
with additions, deletions or substitutions which do not substantially affect the
immunogenic properties of the recombinant polypeptides or polynucleotides.

Compositions, kits and administration

In a further aspect of the invention there are provided compositions comprising a BASB125 polynucleotide and/or a BASB125 polypeptide for administration to a cell or to a multicellular organism.

- 5 The invention also relates to compositions comprising a polynucleotide and/or a polypeptides discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to an individual. Such compositions comprise, for
10 instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one
15 or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

- 20 The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.
- 25 In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

- In a further aspect, the present invention provides for pharmaceutical compositions
30 comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such

as the soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further
5 relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

10 The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic
15 acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.

20 For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course,
25 be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the

attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination,

nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

5 A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

10 A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

15 All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent
20 application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

DEFINITIONS

25 "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known
30 methods, including but not limited to those described in (*Computational Molecular*

- Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heine, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)). BLASTP, BLASTN (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990), and FASTA(Pearson and Lipman Proc. Natl. Acad. Sci. USA 85: 2444-2448 (1988). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman algorithm may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

- Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)
Comparison matrix: BLOSSUM62 from Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)
Gap Penalty: 8
Gap Length Penalty: 2
A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

- Algorithm: Needleman and Wunsch. *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

Available as: The "gap" program from Genetics Computer Group, Madison WI. These
5 are the default parameters for nucleic acid comparisons.

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

- 10 (1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference
15 sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more
20 contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

25
$$n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for
30 the multiplication operator, and wherein any non-integer product of x_n and y is rounded

down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

5

By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain

integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino

acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

"Individual(s)," when used herein with reference to an organism, means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA including single and double-stranded regions.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide.

Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below.

5 A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or
10 inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

15 "Disease(s)" means any disease caused by or related to infection by a bacteria, including, for example, otitis media in infants and children, pneumonia in elderlies, sinusitis, nosocomial infections and invasive diseases, chronic otitis media with hearing loss, fluid accumulation in the middle ear, auditive nerve damage. delayed speech learning, infection
20 of the upper respiratory tract and inflammation of the middle ear.

EXAMPLES:

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1: DNA sequencing of the BASB125 gene from *Moraxella catarrhalis* strain ATCC 43617.

The DNA sequence of the BASB125 gene from the *Moraxella catarrhalis* strain ATCC 43617 (also referred to as strain MC2931) is shown in SEQ ID N0:1. The translation of the BASB125 polynucleotide sequence showed in SEQ ID N0:2.

Example 2: Construction of Plasmid to Express Recombinant BASB125

A: Cloning of BASB125.

The *EcoRI* and *SalI* restriction sites engineered into the MC-Lip17-Fn/t-RI (5'- AGG CAG AGG GAA TTC ATG ATG AAA ATT TTA TAT GTT ACC G -3') [SEQ ID NO:3] forward and MC-Lip17RCh/t-Sal (5'-AGG CAG AGG GTC GAC TTA ATG GTG ATG GTG ATG GTG GCG TGT GGT AGT TGG CAC AAT AAA-3') [SEQ ID NO:4] reverse amplification primers, respectively, permitted directional cloning a PCR product into the *E.coli* expression plasmid pTLZ2 such that a mature BASB125 protein could be expressed as a fusion protein containing a (His)₆ affinity chromatography tag at the C-terminus. The BASB125 PCR product was purified from the amplification reaction using silica gel-based spin columns (QiaGen) according to the manufacturers instructions. To produce the required *EcoRI* and *SalI* termini necessary for cloning, purified PCR product was sequentially digested to completion with *EcoRI* and *SalI* restriction enzymes as recommended by the manufacturer (Life Technologies).

Following the first restriction digestion, the PCR product was purified via spin column as above to remove salts and eluted in sterile water prior to the second enzyme digestion. The digested DNA fragment was again purified using silica gel-based spin columns prior to ligation with the pTLZ2 plasmid.

5

B: Production of Expression Vector.

To prepare the expression plasmid pTLZ2 for ligation, it was similarly digested to completion with both *EcoRI* and *SalI* and then treated with calf intestinal phosphatase (CIP, ~0.02 units / pmole of 5' end, Life Technologies) as directed by the manufacturer to prevent self ligation. An approximately 5-fold molar excess of the digested fragment to the prepared vector was used to program the ligation reaction. A standard ~20 µl ligation reaction (~16°C, ~16 hours), using methods well known in the art, was performed using T4 DNA ligase (~2.0 units / reaction, Life Technologies). An aliquot of the ligation (~5 µl) was used to transform electro-competent JM109 cells according to methods well known in the art. Following a ~2-3 hour outgrowth period at 37°C in ~1.0 ml of LB broth, transformed cells were plated on LB agar plates containing ampicillin (100 µg/ml). Antibiotic was included in the selection. Plates were incubated overnight at 37°C for ~16 hours. Individual ApR colonies were picked with sterile toothpicks and used to "patch" inoculate fresh LB ApR plates as well as a ~1.0 ml LB ApR broth culture. Both the patch plates and the broth culture were incubated overnight at 37°C in either a standard incubator (plates) or a shaking water bath. A whole cell-based PCR analysis was employed to verify that transformants contained the BASB125 DNA insert. Here, the ~1.0 ml overnight LB Ap broth culture was transferred to a 1.5 ml polypropylene tube and the cells collected by centrifugation in a Beckmann microcentrifuge (~3 min., room temperature, ~12,000 X g). The cell pellet was suspended in ~200µl of sterile water and a ~10% aliquot used to program a ~50µl final volume PCR reaction containing both BASB125 forward and reverse amplification primers. Final concentrations of the PCR reaction components were essentially the same as those specified in example 2 except ~5.0 units of *Taq* polymerase was used. The initial 95°C denaturation step was increased to 3 minutes to ensure thermal

disruption of the bacterial cells and liberation of plasmid DNA. An ABI Model 9700 thermal cycler and a 32 cycle, three-step thermal amplification profile, i.e. 95°C. 45sec; 55-58°C, 45sec, 72°C, 1min., were used to amplify the BASB125 fragment from the lysed transformant samples. Following thermal amplification, a ~20µl aliquot of the reaction was analyzed by agarose gel electrophoresis (0.8 % agarose in a Tris-acetate-EDTA (TAE) buffer). DNA fragments were visualized by UV illumination after gel electrophoresis and ethidium bromide staining. A DNA molecular size standard (1 Kb ladder, Life Technologies) was electrophoresed in parallel with the test samples and was used to estimate the size of the PCR products. Transformants that produced the expected size PCR product were identified as strains containing a BASB125 expression construct. Expression plasmid containing strains were then analyzed for the inducible expression of recombinant BASB125.

C: Expression Analysis of PCR-Positive Transformants.

For each PCR-positive transformant identified above, ~5.0 ml of LB broth containing ampicillin (100 µg/ml) was inoculated with cells from the patch plate and grown overnight at 37 °C with shaking (~250 rpm). An aliquot of the overnight seed culture (~1.0 ml) was inoculated into a 125 ml erlenmeyer flask containing ~25 of LB Ap broth and grown at 37 °C with shaking (~250 rpm) until the culture turbidity reached O.D.600 of ~0.5, i.e. mid-log phase (usually about 1.5 - 2.0 hours). At this time approximately half of the culture (~12.5 ml) was transferred to a second 125 ml flask and expression of recombinant BASB125 protein induced by the addition of IPTG (1.0 M stock prepared in sterile water, Sigma) to a final concentration of 1.0 mM. Incubation of both the IPTG-induced and non-induced cultures continued for an additional ~4 hours at 37 °C with shaking. Samples (~1.0 ml) of both induced and non-induced cultures were removed after the induction period and the cells collected by centrifugation in a microcentrifuge at room temperature for ~3 minutes. Individual cell pellets were suspended in ~50µl of sterile water, then mixed with an equal volume of 2X Laemelli SDS-PAGE sample buffer containing 2-mercaptoethanol, and placed in boiling water bath for ~3 min to denature protein. Equal volumes (~15µl) of both the crude IPTG-

induced and the non-induced cell lysates were loaded onto duplicate 12% Tris/glycine polyacrylamide gel (1 mm thick Mini-gels, Novex). The induced and non-induced lysate samples were electrophoresed together with prestained molecular weight markers (SeeBlue, Novex) under conventional conditions using a standard SDS/Tris/glycine running buffer (BioRad). Following electrophoresis, one gel was stained with commassie brilliant blue R250 (BioRad) and then destained to visualize novel BASB125 IPTG-inducible protein(s). The second gel was electroblotted onto a PVDF membrane (0.45 micron pore size, Novex) for ~2 hrs at 4 °C using a BioRad Mini-Protean II blotting apparatus and Towbin's methanol (20 %) transfer buffer. Blocking of the membrane and antibody incubations were performed according to methods well known in the art. A monoclonal anti-RGS (His)3 antibody, followed by a second rabbit anti-mouse antibody conjugated to HRP (QiaGen), was used to confirm the expression and identity of the BASB125 recombinant protein. Visualization of the anti-His antibody reactive pattern was achieved using either an ABT insoluble substrate or using Hyperfilm with the Amersham ECL chemiluminescence system.

Example 3: Production of Recombinant BASB125

Bacterial strain

A recombinant expression strain of *E. coli* JM109 containing a plasmid (pTLZ2) encoding BASB125 from *M. catarrhalis*. was used to produce cell mass for purification of recombinant protein. The expression strain was cultivated on LB agar plates containing 100µg/ml ampicillin ("Ap") to ensure that the pTLZ2 was maintained. For cryopreservation at -80 °C, the strain was propagated in LB broth containing the same concentration of antibiotics then mixed with an equal volume of LB broth containing 30% (w/v) glycerol.

Media

The fermentation medium used for the production of recombinant protein consisted of 2X YT broth (Difco) containing 100µg/ml Ap. Antifoam was added to medium for the fermentor at 0.25 ml/L (Antifoam 204, Sigma). To induce expression of the BASB125

recombinant protein, IPTG (Isopropyl β -D-Thiogalactopyranoside) was added to the fermentor (1 mM, final).

Fermentation

5 A 500-ml erlenmeyer seed flask, containing 50ml working volume, was inoculated with 0.3 ml of rapidly thawed frozen culture, or several colonies from a selective agar plate culture, and incubated for approximately 12 hours at $37 \pm 1^\circ\text{C}$ on a shaking platform at 150rpm (Innova 2100, New Brunswick Scientific). This seed culture was then used to inoculate a 5-L working volume fermentor containing 2X YT broth and both Ap
10 antibiotics. The fermentor (Bioflo 3000, New Brunswick Scientific) was operated at $37 \pm 1^\circ\text{C}$, 0.2 - 0.4 VVM air sparge, 250 rpm in Rushton impellers. The pH was not controlled in either the flask seed culture or the fermentor. During fermentation, the pH ranged 6.5 to 7.3 in the fermentor. IPTG (1.0 M stock, prepared in sterile water) was added to the fermentor when the culture reached mid-log of growth (~ 0.7 O.D.600
15 units). Cells were induced for 2 - 4 hours then harvested by centrifugation using either a 28RS Heraeus (Sepatech) or RC5C superspeed centrifuge (Sorvall Instruments). Cell paste was stored at -20°C until processed.

Chemicals and Materials

20 Imidazole, guanidine hydrochloride, Tris (hydroxymethyl), and EDTA (ethylene-diamine tetraacetic acid) biotechnology grade or better were all obtained from Ameresco Chemical, Solon, Ohio. Triton X-100 (t-Octylphenoxypolyethoxy-ethanol), Triton X-114, and sodium phosphate, monobasic were reagent grade or better and obtained from Sigma Chemical Company, St. Louis, Missouri. Glacial acetic acid and
25 hydrochloric acid were obtained from Mallincrodt Baker Inc., Phillipsburg, New Jersey. Methanol was obtained from Fisher Scientific, Fairlawn, New Jersey. Pefabloc®SC (4-(2-Aminoethyl)-benzenesulfonylfluoride), Complete protease inhibitor cocktail tablets, and PMSF (phenylmethyl-sulfonylfluoride) were obtained from Roche Diagnostics Corporation, Indianapolis, Indiana. Bestatin, Pepstatin A, and E-64 protease inhibitor
30 were obtained from Calbiochem, LaJolla, California. Dulbecco's Phosphate Buffered Saline(1x PBS) was obtained from Quality Biological, Inc., Gaithersburg, Maryland.

Dulbecco's Phosphate Buffered Saline (10x PBS) was obtained from BioWhittaker, Walkersville, Maryland. Penta-His Antibody, BSA free was obtained from QiaGen, Valencia, California. Peroxidase-conjugated AffiniPure Goat Anti-mouse IgG was obtained from Jackson Immuno Research, West Grove, Penn. AEC single solution was
5 obtained from Zymed, South San Francisco, California. All other chemicals were reagent grade or better.

Ni-Chelating Sepharose Fast Flow resin was obtained from Pharmacia., Sweden, California. Precast Tris-Glycine 4-20% and 10-20% polyacrylamide gels, all running buffers and solutions, SeeBlue Pre-Stained Standards, MultiMark Multi-Colored
10 Standards and PVDF transfer membranes were obtained from Novex, San Diego, California. SDS-PAGE Silver Stain kits were obtained from Daiichi Pure Chemicals Company Limited, Tokyo, Japan. Coomassie Stain Solution was obtained from Bio-Rad Laboratories, Hercules, California. Acrodisc® PF 0.2 µm syringe filters were obtained from Pall Gelman Sciences, Ann Arbor, Michigan. GD/X 25mm disposable
15 syringe filters were obtained from Whatman Inc., Clifton, New Jersey. Dialysis tubing 8,000 MWCO was obtained from BioDesign Inc. Od New York, Carmal New York. BCA Protein Assay Reagents and Snake Skin dialysis tubing 3,500 MWCO were obtained from Pierce Chemical Co., Rockford, Illinois.

20 **Example 4: Purification of recombinant BASB125 from *E. coli***

Extraction Purification

Cell paste was thawed at room temperature for 30 to 60 minutes. Following disruption of the cell paste, the extract was partitioned with ice cold PBS containing 1% TritonX-114.

25 Cell debris were removed, the extract was warmed to 37° C and the phases partitioned by centrifugation.

The fraction of interest was then passed over a nickel-Chelating Sepharose Fast Flow resin equilibrated in PBS (pH 7.5) containing 10 % glycerol and 0.05 % Triton X100. The protein was eluted in the same buffer containing 200 mM Imidazole. The fraction
30 containing the eluted protein was diluted with four volumes of 50 mM Tris buffer (pH

7.5) containing 2 mM EDTA, 10 mM sodium chloride and 0.05% Triton X100 and run through a DEAE-Sepharose FF resin. The flow through was collected and concentrated on a 3 kDa cut-off stir cell, then dialysed against PBS (pH 7.4) containing 0.1% Triton X100.

5

As shown in figure 1-A, purified BASB125 protein appeared in SDS-PAGE analysis as 3 bands migrating at around 14 and 6 kDa (estimated relative molecular mass). Purity was estimated around to 50 %. Only one was reactive against a mouse monoclonal antibody raised against the 6-Histidine motif (figure 1-B).

10

Biochemical Characterizations

SDS-PAGE and Western Blot Analysis

The recombinant purified BASB125 protein was resolved on 4-20 % polyacrylamide gels and electrophoretically transferred to PVDF membranes at 100 V for 1 hour as previously described (Thebaine et al. 1979, Proc. Natl. Acad. Sci. USA 76:4350-4354). The PVDF membranes were then pretreated with 25 ml of Dulbecco's phosphate buffered saline containing 5 % non-fat dry milk. All subsequent incubations were carried out using this pretreatment buffer.

20

PVDF membranes were incubated with 25 ml of a 1:500 dilution of preimmune or immune serum or anti-His immune serum for 1 hour at room temperature. PVDF membranes were then washed twice with wash buffer (20 mM Tris buffer, pH 7.5, containing 150 mM sodium chloride and 0.05 % Tween-20). PVDF membranes were incubated with 25 ml of a 1:5000 dilution of peroxidase-labeled goat anti-rabbit or anti-mouse IgG (Jackson ImmunoResearch Laboratories, West Grove, PA) for 30 minutes at room temperature. PVDF membranes were then washed 4 times with wash buffer, and were developed with 3-amino-9-ethylcarbazole and urea peroxide as supplied by Zymed (San Francisco, CA) for 10 minutes each.

30

Example 5: Production of Antisera to Recombinant BASB125

Polyvalent antisera directed against the BASB125 protein were generated by vaccinating two rabbits with the purified recombinant BASB125 protein. Each animal is given a total of three immunizations subcutaneously of about 10µg BASB125 protein per injection at approximately 21 days interval. Animals were bled prior to the first immunization ("pre-bleed") and on days 49 and 56.

Anti-BASB125 protein titers were measured by an ELISA using purified recombinant BASB125 protein (4 µg/well). The titre is defined as mid-point titers calculated by 4-parameter logistic model using the XL Fit software. The titers obtained post immunisation were 1:600 for the rabbits sera.

The antisera were used as the first antibody to identify the protein in a western blot. The western-blot shows the presence of anti-BASB125 antibody in the sera of immunized animals (figure 2, lanes indicated by an arrow).

Example 6: Immunological characterization: Surface exposure of BASB125

Anti-BASB125 protein titers were determined by an ELISA using formalin-killed whole cells of *Moraxella catarrhalis* strains 358,216,2926 (20 µg/well) The titre is defined as mid-point titers calculated by 4-parameter logistic model using the SoftMax Pro software.

Titers observed with the rabbit immune sera (1:396,1:285,1:848 respectively) demonstrate that the BASB125 protein is detected at the surface of *M. catarrhalis* cells.

Example 7. Immunological Characterisation: Western Blot Analysis

Several strains of *M. catarrhalis* including ATCC 43617, as well as clinical isolates from various geographic regions, are grown on Muller Hinton agar plates for 24 hours at 36°C. Several colonies are used to inoculate broth. Cultures are grown until the A620 is approximately 0.6 and cells are collected by centrifugation. Cells are then

concentrated and solubilized in PAGE sample buffer. The solubilized cells are then resolved on 4-20% polyacrylamide gels and the separated proteins are electrophoretically transferred to PVDF membranes. The PVDF membranes are then pretreated with saturation buffer. All subsequent incubations are carried out using this pretreatment buffer.

PVDF membranes are incubated with preimmune serum or rabbit or mouse immune seru. PVDF membranes are then washed.

PVDF membranes are incubated with biotin-labeled sheep anti-rabbit or mouse Ig.

PVDF membranes are then washed 3 times with wash buffer, and incubated with streptavidin-peroxydase. PVDF membranes are then washed 3 times with wash buffer and developed with 4-chloro-1-naphtol.

Detection of a protein corresponding to BASB125 expected molecular weight that is reactive with the antisera is used to demonstrate that this protein is produced by and conserved in all *Moraxella* strains tested.

Example 8: Immunological characterization: Bactericidal Activity

Complement-mediated cytotoxic activity of anti-BASB125 antibodies is examined to determine the vaccine potential of BASB125 protein antiserum that is prepared as described above. The activities of the pre-immune serum and the anti-BASB125 antiserum in mediating complement killing of *M. catarrhalis* are examined.

Strains of *M. catarrhalis* are grown on plates. Several colonies are added to liquid medium. Cultures are grown and collected until the A620 is approximately 0.4. After one wash step, the pellet is suspended and diluted.

Preimmune sera and the anti-BASB125 sera is deposited into the first well of a 96-wells plate and serial dilutions are deposited in the other wells of the same line. Live diluted *M. catarrhalis* is subsequently added and the mixture is incubated. Complement is added into each well at a working dilution defined beforehand in a toxicity assay.

Each test includes a complement control (wells without serum containing active or inactivated complement source), a positive control (wells containing serum with a know titer of bactericidal antibodies), a culture control (wells without serum and complement) and a serum control (wells without complement).

5

Bactericidal activity of rabbit or mice antiserum (50% killing of homologous strain) is measured.

Example 9: Presence of Antibody to BASB125 in Human Convalescent Sera

10 Western blot analysis of purified recombinant BASB125 is performed as described in Example 7 above, except that a pool of human sera from children infected by *M. catarrhalis* is used as the first antibody preparation. This is used to show that antisera from naturally infected individuals react to the purified recombinant protein.

15 **Example 10: Efficacy of BASB125 vaccine: enhancement of lung clearance of *M. catarrhalis* in mice.**

This mouse model is based on the analysis of the lung invasion by *M. catarrhalis* following a standard intranasal challenge to vaccinated mice.

20 Groups of mice are immunized with BASB125 vaccine. After the booster, the mice are challenged by instillation of bacterial suspension into the nostril under anaesthesia. Mice are killed between 30 minutes and 24 hours after challenge and the lungs are removed aseptically and homogenized individually. The log10 weighted mean number of CFU/lung is determined by counting the colonies grown on agar plates after plating
25 of dilutions of the homogenate. The arithmetic mean of the log10 weighted mean number of CFU/lung and the standard deviations are calculated for each group. Results are analysed statistically.

In this experiment groups of mice are immunized either with BASB125 or with a killed
30 whole cells (kwc) preparation of *M. catarrhalis* or sham immunized.

This is used to show that the kwc preparation and the BASB125 vaccine induce significant lung clearance as compared to the control group.

5 **Example 11: Inhibition of *M. catarrhalis* adhesion onto cells by anti-BASB125 antiserum.**

This assay measures the capacity of anti BASB125 sera to inhibit the adhesion of *Moraxella* bacteria to epithelial cells. This activity could prevent colonization of f.i. the nasopharynx by *Moraxella*.

- 10 One volume of bacteria is incubated on ice with one volume of pre-immune or anti-BASB125 immune serum dilution. This mixture is subsequently added in the wells of a 24 well plate containing a confluent cells culture that is washed once with culture medium to remove traces of antibiotic. The plate is centrifuged and incubated.
- 15 Each well is then gently washed. After the last wash, sodium glycocholate is added to the wells. After incubation, the cell layer is scraped and homogenised. Dilutions of the homogenate are plated on agar plates and incubated. The number of colonies on each plate is counted and the number of bacteria present in each well calculated.

This is used to show that bacteria incubated with anti-BASB125 antiserum are inhibited in their adherence capacity to the Hep-2 cells.

20

25

Deposited materials

5 A deposit containing a *Moraxella catarrhalis* Catlin strain has been deposited with the American Type Culture Collection (herein "ATCC") on June 21, 1997 and assigned deposit number 43617. The deposit was described as *Branhamella catarrhalis* (Frosch and Kolle) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *M. catarrhalis* isolate obtained from a transtracheal aspirate of a coal miner with chronic bronchitis. The deposit is described in Antimicrob. Agents Chemother. 21: 506-508 (1982).

10 The *Moraxella catarrhalis* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains a full length BASB125 gene.

15 A deposit of the vector pMC-ORF1/2 consisting of *Moraxella catarrhalis* DNA inserted in pQE30 has been deposited with the American Type Culture Collection (ATCC) on February 12th 1999 and assigned deposit number 207118.

20 The sequence of the polynucleotides contained in the deposited strain / clone, as well as the amino acid sequence of any polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

25 The deposit of the deposited strains have been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The deposited strains will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strains are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 62 lines 5-29.	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet <input type="checkbox"/>
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 UNIVERSITY BLVD, MANASSAS, VIRGINIA 20110-2209, UNITED STATES OF AMERICA	
Date of deposit 21 June 1997 and 12 February 1999	Accession Number 43617 and 207118
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations where a European Patent is sought, a sample of the deposited microorganisms will be made available until the publication of the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn, only by issue of such a sample to an expert nominated by the person requesting the sample	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input checked="checked" type="checkbox"/>	This sheet was received with the international application
Authorized officer N. Mailward	

For International Bureau use only	
<input type="checkbox"/>	This sheet was received by the International Bureau on:
Authorized officer	

Form PCT/RO/134 (July 1992)

SEQUENCE INFORMATION

BASB125 Polynucleotide and Polypeptide Sequences

SEQ ID NO:1

5 *Moraxella catarrhalis* BASB125 polynucleotide sequence from strain ATCC43617

ATGATGAAAATTTTATATGTTACCGCCACACTAATGACTGCCTTTACCCTAGCAAGCTGT
GCAAGTACCCCTGAAAGCAATCCAAAAACAGTTCTGCTAATTTAACCACCAGCTTAATC
AAACATGCAGTCAAACAAACCTGTCAAACCCAGCTGACTGGTCATCAGTACTGGAAAATT
GCCGCCATGAAATTGTCTTCAGAATCCAAAGCCAAGATTTAGAAACGGCATGCGGCTGT
10 GTGGCTGATAAAGCACCTGAAGCCGTGAGCTTGACCGAGCTTACCACAGCTGCCATTAAT
CCAAATGCACGCACTGAAGTTGCCCAAAAATTGTGCGACACTCGCTTAAACCTTGCATG
CTAGAGACCGTCAATGCTTTTATTGTGCCAACTACCACACGCTAA

SEQ ID NO:2

15 *Moraxella catarrhalis* BASB125 polypeptide sequence deduced from the
polynucleotide of SEQ ID NO:1

MMKILYVTATLMTAFTLASCASTPESNPKNSSANLTTSLIKHAVKQTCQTOLTGHQYWKI
AAMKLSSSEKAKISETACGCVADKAPEAVSLTELTTAAINPNARTEVAQKIVRHSLKPCM
20 LETVNAFIVPTTTR

SEQ ID NO:3

AGG CAG AGG GAA TTC ATG ATG AAA ATT TTA TAT GTT ACC G

25 SEQ ID NO:4

AGG CAG AGG GTC GAC TTA ATG GTG ATG GTG ATG GTG GCG TGT GGT
AGT TGG CAC AAT AAA

CLAIMS:

1. An isolated polypeptide comprising an amino acid sequence which has at least 85% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID
5 NO:2.

2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of
10 SEQ ID NO:2.

3. The polypeptide as claimed in claim 1 comprising the amino acid sequence of SEQ ID
15 NO:2.

4. An isolated polypeptide of SEQ ID NO:2.

5. An immunogenic fragment of the polypeptide as claimed in any one of claims 1 to 4 in which the immunogenic activity of said immunogenic fragment is substantially the same as the polypeptide of SEQ ID NO:2.

6. A polypeptide as claimed in any of claims 1 to 5 wherein said polypeptide is part of a larger fusion protein.

7. An isolated polynucleotide encoding a polypeptide as claimed in any of claims 1 to 6.

8. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least 85% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2; or a nucleotide sequence complementary to said isolated polynucleotide.

9. An isolated polynucleotide comprising a nucleotide sequence that has at least 85% identity to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2 over the entire coding region; or a nucleotide sequence complementary to said isolated polynucleotide.

10. An isolated polynucleotide which comprises a nucleotide sequence which has at least 85% identity to that of SEQ ID NO:1 over the entire length of SEQ ID NO:1; or a nucleotide sequence complementary to said isolated polynucleotide.

5

11. The isolated polynucleotide as claimed in any one of claims 7 to 10 in which the identity is at least 95% to SEQ ID NO:1.

10

12. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2.

13. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1.

15

14. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof.

20

15. An expression vector or a recombinant live microorganism comprising an isolated polynucleotide according to any one of claims 7 - 14.

25

16. A host cell comprising the expression vector of claim 15 or a subcellular fraction or a membrane of said host cell expressing an isolated polypeptide comprising an amino acid sequence that has at least 85% identity to the amino acid sequence of SEQ ID NO:2.

30

17. A process for producing a polypeptide of claims 1 to 6 comprising culturing a host cell of claim 16 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.

18. A process for expressing a polynucleotide of any one of claims 7 – 14 comprising transforming a host cell with the expression vector comprising at least one of said polynucleotides and culturing said host cell under conditions sufficient for expression of any one of said polynucleotides.

5

19. A vaccine composition comprising an effective amount of the polypeptide of any one of claims 1 to 6 and a pharmaceutically acceptable carrier.

10

20. A vaccine composition comprising an effective amount of the polynucleotide of any one of claims 7 to 14 and a pharmaceutically effective carrier.

21. The vaccine composition according to either one of claims 19 or 20 wherein said composition comprises at least one other *Moraxella catarrhalis* antigen.

15

22. An antibody immunospecific for the polypeptide or immunological fragment as claimed in any one of claims 1 to 6.

20

23. A method of diagnosing a *Moraxella catarrhalis* infection, comprising identifying a polypeptide as claimed in any one of claims 1 - 6, or an antibody that is immunospecific for said polypeptide, present within a biological sample from an animal suspected of having such an infection.

25

24. Use of a composition comprising an immunologically effective amount of a polypeptide as claimed in any one of claims 1 – 6 in the preparation of a medicament for use in generating an immune response in an animal.

30

25. Use of a composition comprising an immunologically effective amount of a polynucleotide as claimed in any one of claims 7 - 14 in the preparation of a medicament for use in generating an immune response in an animal.

26. A therapeutic composition useful in treating humans with *Moraxella catarrhalis* disease comprising at least one antibody directed against the polypeptide of claims 1 – 6 and a suitable pharmaceutical carrier.

Figure 1-A : Coomassie stained SDS-polyacrylamide gel of purified BASB125

Figure 1-B: Western-blotting of purified BASB125 (anti-His antibody).

Fig. 1-A




Fig. 1-B





Figure 2 : Detection of BASB125 with rabbit antisera (lanes indicated by an arrow).



DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

NOVEL COMPOUNDS the specification of which (check one)

☐ is attached hereto.

☒ was filed on 27 July 2000 as Serial No. PCT/EP00/07291
and was amended on (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or Inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Number	Country	Filing Date	Priority Claimed
9918041.6	Great Britain	30 July 1999	Yes

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

Application Number Filing Date

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each

of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.


Serial No.	Filing Date	Status
------------	-------------	--------

I hereby appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

Customer Number 25,308

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-60 Full Name of Inventor: Joelle THONNARD

Inventor's Signature: 

Date: 4 December 2007

Residence: Rixensart, Belgium BEL

Citizenship: Belgian

Post Office Address: SmithKline Beecham Corporation
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939

SEQUENCE LISTING

<110> Joelle THONNARD

<120> Novel Compounds

<130> BM45402

<150> 9918041.6

<151> 1999-07-30

<160> 4

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 405

<212> DNA

<213> Moraxella catarrhalis

<400> 1

atgatgaaaa	ttttatatgt	taccgccaca	ctaagtactg	cctttaccct	agcaagctgt	60
gcaagtaccc	ctgaaagcaa	tccaaaaaac	agttctgcta	atttaaccac	cagcttaatc	120
aaacatgcag	tcaaacaaac	ctgtcaaacc	cagctgactg	gtcatcagta	ctggaaaatt	180
gccgccatga	aattgtcttc	agaatccaaa	gccaaagattt	cagaaaacggc	atgcggctgt	240
gtggctgata	aagcacctga	agccgtcagc	ttgaccgagc	ttaccacagc	tgccattaat	300
ccaaatgcac	gactgaagt	tgcccaaaaa	attgtgcgac	actcgcttaa	accttgcattg	360
ctagagaccg	tcaatgcttt	tattgtgcc	actaccacac	gctaa		405

<210> 2

<211> 134

<212> PRT

<213> Moraxella catarrhalis

<400> 2

Met	Met	Lys	Ile	Leu	Tyr	Val	Thr	Ala	Thr	Leu	Met	Thr	Ala	Phe	Thr
1				5					10					15	
Leu	Ala	Ser	Cys	Ala	Ser	Thr	Pro	Glu	Ser	Asn	Pro	Lys	Asn	Ser	Ser
		20						25					30		
Ala	Asn	Leu	Thr	Thr	Ser	Leu	Ile	Lys	His	Ala	Val	Lys	Gln	Thr	Cys
		35				40						45			
Gln	Thr	Gln	Leu	Thr	Gly	His	Gln	Tyr	Trp	Lys	Ile	Ala	Ala	Met	Lys
	50				55					60					
Leu	Ser	Ser	Glu	Ser	Lys	Ala	Lys	Ile	Ser	Glu	Thr	Ala	Cys	Gly	Cys
65				70					75					80	
Val	Ala	Asp	Lys	Ala	Pro	Glu	Ala	Val	Ser	Leu	Thr	Glu	Leu	Thr	Thr
			85					90					95		
Ala	Ala	Ile	Asn	Pro	Asn	Ala	Arg	Thr	Glu	Val	Ala	Gln	Lys	Ile	Val
			100					105					110		
Arg	His	Ser	Leu	Lys	Pro	Cys	Met	Leu	Glu	Thr	Val	Asn	Ala	Phe	Ile
		115					120								
Val	Pro	Thr	Thr	Thr	Arg										


Parameter	Unit	Value
Mean	mm	1.0
Standard deviation	mm	0.2
Minimum	mm	0.5
Maximum	mm	1.5
Range	mm	1.0
Median	mm	1.0
Mode	mm	1.0
Skewness		0.0
Kurtosis		0.0
Correlation coefficient		1.0
Regression equation		$y = x$
Intercept	mm	0.0
Slope		1.0
Adjusted R-squared		1.0
F-statistic		1.0
P-value		0.0
Confidence interval	mm	0.8 to 1.2
95% confidence interval	mm	0.7 to 1.3
90% confidence interval	mm	0.6 to 1.4
80% confidence interval	mm	0.5 to 1.5
70% confidence interval	mm	0.4 to 1.6
60% confidence interval	mm	0.3 to 1.7
50% confidence interval	mm	0.2 to 1.8
40% confidence interval	mm	0.1 to 1.9
30% confidence interval	mm	0.0 to 2.0
20% confidence interval	mm	-0.1 to 2.1
10% confidence interval	mm	-0.2 to 2.2
5% confidence interval	mm	-0.3 to 2.3
1% confidence interval	mm	-0.4 to 2.4
0.5% confidence interval	mm	-0.5 to 2.5
0.1% confidence interval	mm	-0.6 to 2.6
0.05% confidence interval	mm	-0.7 to 2.7
0.01% confidence interval	mm	-0.8 to 2.8
0.005% confidence interval	mm	-0.9 to 2.9
0.001% confidence interval	mm	-1.0 to 3.0
0.0005% confidence interval	mm	-1.1 to 3.1
0.0001% confidence interval	mm	-1.2 to 3.2
0.00005% confidence interval	mm	-1.3 to 3.3
0.00001% confidence interval	mm	-1.4 to 3.4
0.000005% confidence interval	mm	-1.5 to 3.5
0.000001% confidence interval	mm	-1.6 to 3.6
0.0000005% confidence interval	mm	-1.7 to 3.7
0.0000001% confidence interval	mm	-1.8 to 3.8
0.00000005% confidence interval	mm	-1.9 to 3.9
0.00000001% confidence interval	mm	-2.0 to 4.0
0.000000005% confidence interval	mm	-2.1 to 4.1
0.000000001% confidence interval	mm	-2.2 to 4.2
0.0000000005% confidence interval	mm	-2.3 to 4.3
0.0000000001% confidence interval	mm	-2.4 to 4.4
0.00000000005% confidence interval	mm	-2.5 to 4.5
0.00000000001% confidence interval	mm	-2.6 to 4.6
0.000000000005% confidence interval	mm	-2.7 to 4.7
0.000000000001% confidence interval	mm	-2.8 to 4.8
0.0000000000005% confidence interval	mm	-2.9 to 4.9
0.0000000000001% confidence interval	mm	-3.0 to 5.0
0.00000000000005% confidence interval	mm	-3.1 to 5.1
0.00000000000001% confidence interval	mm	-3.2 to 5.2
0.000000000000005% confidence interval	mm	-3.3 to 5.3
0.000000000000001% confidence interval	mm	-3.4 to 5.4
0.0000000000000005% confidence interval	mm	-3.5 to 5.5
0.0000000000000001% confidence interval	mm	-3.6 to 5.6
0.00000000000000005% confidence interval	mm	-3.7 to 5.7
0.00000000000000001% confidence interval	mm	-3.8 to 5.8
0.000000000000000005% confidence interval	mm	-3.9 to 5.9
0.000000000000000001% confidence interval	mm	-4.0 to 6.0
0.0000000000000000005% confidence interval	mm	-4.1 to 6.1
0.0000000000000000001% confidence interval	mm	-4.2 to 6.2
0.00000000000000000005% confidence interval	mm	-4.3 to 6.3
0.00000000000000000001% confidence interval	mm	-4.4 to 6.4
0.000000000000000000005% confidence interval	mm	-4.5 to 6.5
0.000000000000000000001% confidence interval	mm	-4.6 to 6.6
0.0000000000000000000005% confidence interval	mm	-4.7 to 6.7
0.0000000000000000000001% confidence interval	mm	-4.8 to 6.8
0.00000000000000000000005% confidence interval	mm	-4.9 to 6.9
0.00000000000000000000001% confidence interval	mm	-5.0 to 7.0
0.000000000000000000000005% confidence interval	mm	-5.1 to 7.1
0.000000000000000000000001% confidence interval	mm	-5.2 to 7.2
0.0000000000000000000000005% confidence interval	mm	-5.3 to 7.3
0.0000000000000000000000001% confidence interval	mm	-5.4 to 7.4
0.00000000000000000000000005% confidence interval	mm	-5.5 to 7.5
0.00000000000000000000000001% confidence interval	mm	-5.6 to 7.6
0.000000000000000000000000005% confidence interval	mm	-5.7 to 7.7
0.000000000000000000000000001% confidence interval	mm	-5.8 to 7.8
0.0000000000000000000000000005% confidence interval	mm	-5.9 to 7.9
0.0000000000000000000000000001% confidence interval	mm	-6.0 to 8.0
0.00000000000000000000000000005% confidence interval	mm	-6.1 to 8.1
0		

```
<220>
<223> primer
```

```
<210> 4
<211> 60
<212> DNA
<213> Artificial Sequence
```

<400> 4
aggcagaggg tgcacttaat ggtgatggtg atggtggcgt gtggtagtgtg gcacaataaa 60

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



SCANNED, # 12

Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ *Scanned copy is best available.*

- Only 2 sheet of drawings not 3.